

Techniques for one-loop tensor integrals in many-particle processes

A. Denner^a and S. Dittmaier^b

^aPaul Scherrer Institut, Würenlingen und Villigen, CH-5232 Villigen PSI, Switzerland

^bMax-Planck-Institut für Physik (Werner-Heisenberg-Institut), D-80805 München, Germany

We briefly sketch the methods for a numerically stable evaluation of tensor one-loop integrals that have been used in the calculation of the complete electroweak one-loop corrections to $e^+e^- \rightarrow 4$ fermions. In particular, the improvement of the new methods over the conventional Passarino–Veltman reduction is illustrated for some 4-point integrals in the delicate limits of small Gram (and other kinematical) determinants.

1. INTRODUCTION

At the LHC and ILC, many interesting processes involve more than four external particles. Such many-particle reactions often proceed via resonances that subsequently decay, or they represent a background to resonance processes. A thorough description of such processes requires the evaluation of strong and electroweak radiative corrections at least in next-to-leading order. The most complicated part in such calculations concerns the numerically stable evaluation of the one-loop tensor integrals of the virtual corrections.

For processes with up to four external particles the classical Passarino–Veltman (PV) reduction [1], which recursively reduces tensor to scalar integrals, is sufficient in practically all cases. This scheme, however, involves Gram determinants in the denominator, which spoil the numerical stability if they become small. With up to four external particles this happens only near the edge of phase space (forward scattering, thresholds). With more than four external particles, Gram determinants also vanish within phase space, and methods are needed where Gram determinants can be small but still non-zero. Several solutions to this problem have been proposed in recent years, but not many of them have proven their performance in complicated applications yet. For references and descriptions of some methods alternative to ours, we refer to Refs. [2,3].

In this article we briefly describe the methods used in the calculation of the complete electroweak $\mathcal{O}(\alpha)$ corrections to $e^+e^- \rightarrow 4$ fermions

[4], which constitutes the first established one-loop result for a $2 \rightarrow 4$ particle reaction. All relevant formulas can be found in Refs. [2,5], here we only sketch their structure. Moreover, we illustrate the improvement of the new methods over the conventional PV reduction for some 4-point integrals in delicate kinematical limits.

2. THE GENERAL CONCEPT

One-loop tensor integrals can be naturally grouped into three categories, which we have treated in completely different ways:

- (i) For *1- and 2-point integrals* of arbitrary tensor rank, numerically stable analytical expressions are presented in Ref. [2] (see also Ref. [1]).
- (ii) For *3- and 4-point tensor integrals*, PV reduction [1] is applied for “regular” phase-space points where Gram determinants are not too small. For the remaining problematic cases special reduction techniques have been developed [2].

One of the techniques replaces the standard scalar integral by a specific tensor coefficient that can be safely evaluated numerically and reduces the remaining tensor coefficients as well as the standard scalar integral to the new basis integrals. In this scheme no dangerous inverse Gram determinants occur, but inverse modified Cayley determinants instead. The procedure is related to the fully numerical method described in Ref. [6].

In a second class of techniques, the tensor coefficients are iteratively deduced up to terms that

are systematically suppressed by small Gram or other kinematical determinants in specific kinematical configurations. The numerical accuracy can be systematically improved upon including higher tensor ranks. A similar idea, where tensor coefficients are iteratively determined from higher-rank tensors has been described in Ref. [7] for the massless case.

These methods are sketched in more detail and illustrated in 4-point examples below.

(iii) For *5- and 6-point integrals*, direct reductions to 5- and 4-point integrals, respectively, are possible owing to the four-dimensionality of space-time. For scalar integrals such a reduction was already derived in the 1960s [10]. In Refs. [2,5] we follow basically the same strategy to reduce tensor integrals, which has the advantage that no inverse Gram determinants appear in the reduction. Instead modified Cayley determinants occur in the denominator, but we did not find numerical problems with these factors. A reduction similar to ours has been proposed in Ref. [8].

We would like to stress two important features of our approach.

(i) The methods are valid for massive and massless cases. The formulas given in Refs. [2,5] are valid without modifications if IR divergences are regularized with mass parameters or dimensionally.¹ Finite masses can be either real or complex.

(ii) The in/out structure of the methods is the same as for conventional PV reduction, i.e. no specific algebraic manipulations are needed in applications. Therefore, the whole method can be (and in fact is) organized as a numerical library for scalar integrals and tensor coefficients.

We conclude this overview with some comments resulting from our experience collected in the treatment of a full $2 \rightarrow 4$ scattering reaction.

(i) For a specific point in a multi-particle (multi-parameter) phase space it is highly non-trivial to figure out which of the various methods is the most precise. It seems hopeless to split the phase space into regions that are dedicated to a given method. Therefore, we estimate the accuracy for the different methods at each phase-space

¹For the method of Ref. [5], this has been shown in Ref. [9].

point and take the variant promising the highest precision. In one approach, we estimate the number of valid digits based on the expected accuracy of the expansions and possible numerical cancellations before the evaluation of the coefficients. In a second approach, the accuracy of the PV method is estimated by checking symmetries and sum rules for the coefficients. If the estimated precision is not satisfactory, the seminumerical method is taken; if this is still not satisfactory, an expansion is used.

(ii) In a complicated phase space it may happen that none of the various methods is perfect or good in some exceptional situations. Usually the corresponding events do not significantly contribute to cross sections. This issue can only be fathomed in actual applications. To be on the safe side, we employ the two independent “rescue systems” with different advantages and limitations.

(iii) In view of this, figures as shown below are nice illustrations, but should always be taken with a grain of salt. No matter how many of such figures are shown, they will never be exhaustive, so that no quantitative conclusions on the overall precision of methods can be drawn.

In summary, the performance (speed, stability, etc.) of any method can only be estimated and proven in non-trivial and realistic applications.

3. 4-POINT INTEGRALS FOR SMALL GRAM DETERMINANTS

In the following we sketch the methods of Ref. [2] for the 4-point tensor integrals of rank P ,

$$D^{\mu_1 \dots \mu_P} = \frac{(2\pi\mu)^{4-D}}{i\pi^2} \int d^D q \frac{q^{\mu_1} \dots q^{\mu_P}}{N_0 N_1 N_2 N_3},$$

$$N_k = (q + p_k)^2 - m_k^2 + i\epsilon, \quad p_0 = 0, \quad (1)$$

which are decomposed into covariants as follows,

$$D^\mu = \sum_{i_1=1}^3 p_{i_1}^\mu D_{i_1},$$

$$D^{\mu\nu} = \sum_{i_1, i_2=1}^3 p_{i_1}^\mu p_{i_2}^\nu D_{i_1 i_2} + g^{\mu\nu} D_{00},$$

$$D^{\mu\nu\rho} = \sum_{i_1, i_2, i_3=1}^3 p_{i_1}^\mu p_{i_2}^\nu p_{i_3}^\rho D_{i_1 i_2 i_3}$$

$$+ \sum_{i_1=1}^3 (g^{\mu\nu} p_{i_1}^\rho + g^{\nu\rho} p_{i_1}^\mu + g^{\rho\mu} p_{i_1}^\nu) D_{00i_1}, \quad (2)$$

and so on for higher rank. The conventional PV reduction [1] recursively expresses the tensor coefficients $D_{i_1 \dots i_P}$ in terms of the scalar 4-point integral D_0 and 3-point integrals of lower rank. To this end, the defining tensor integrals are contracted with an external momentum $p_k^{\mu_1}$ or with the metric tensor $g^{\mu_1 \mu_2}$, leading to the factors $p_k q$ or q^2 in the numerator, respectively. Rewriting these factors according to

$$\begin{aligned} 2p_k q &= N_k - N_0 - f_k, & f_k &= p_k^2 - m_k^2 + m_0^2, \\ q^2 &= N_0 + m_0^2, \end{aligned} \quad (3)$$

the N_i terms can be used to cancel propagator denominators. This procedure yields the basic PV relations that connect contracted 4-point tensor integrals of rank P with lower-rank 4- and 3-point integrals. The PV solution expresses the rank P 4-point coefficients in terms of lower-rank 4- and 3-point coefficients. In each step $P \rightarrow (P-1)$ the inverse of the Gram matrix

$$Z = \begin{pmatrix} 2p_1 p_1 & 2p_1 p_2 & 2p_1 p_3 \\ 2p_2 p_1 & 2p_2 p_2 & 2p_2 p_3 \\ 2p_3 p_1 & 2p_3 p_2 & 2p_3 p_3 \end{pmatrix} \quad (4)$$

occurs, which causes the above-mentioned numerical problems if the determinant $|Z|$ becomes small. In our alternative methods also the matrix

$$X = \left(\begin{array}{c|ccc} 2m_0^2 & f_1 & f_2 & f_3 \\ \hline f_1 & & & \\ f_2 & & Z & \\ f_3 & & & \end{array} \right), \quad (5)$$

and its inverse appear. The vanishing of the modified Cayley determinant $|X|$ corresponds to necessary conditions for true (Landau) singularities in a Feynman diagram.

We adopt the conventions that indices $\hat{i}$ with a hat should be omitted and that $\tilde{Z}_{ij}$ and $\tilde{X}_{ij}$ denote the minors (i.e. determinants of submatrices where row i and column j are discarded) of the matrices Z and X , respectively. Similarly $\tilde{Z}_{ij,kl}$ denotes generalized minors where two rows and two columns are discarded in Z . For the precise definitions, full formulas, and detailed descriptions of algorithms we refer to Ref. [2]. Here we only highlight the general structure.

3.1. Seminumerical method for small $|Z|$

The PV relations can be arranged as follows,

$$X \begin{pmatrix} D_{i_2 \dots i_P} \\ D_{1i_2 \dots i_P} \\ D_{2i_2 \dots i_P} \\ D_{3i_2 \dots i_P} \end{pmatrix} = -2 \begin{pmatrix} \text{const.} \times D_{00i_2 \dots i_P} \\ \sum_{r=2}^P \delta_{1i_r} D_{00i_2 \dots \hat{i}_r \dots i_P} \\ \sum_{r=2}^P \delta_{2i_r} D_{00i_2 \dots \hat{i}_r \dots i_P} \\ \sum_{r=2}^P \delta_{3i_r} D_{00i_2 \dots \hat{i}_r \dots i_P} \end{pmatrix} + 3\text{-point integrals}, \quad (6)$$

which expresses tensor coefficients of rank P and $(P-1)$ in terms of coefficients of rank P and $(P+1)$ that involve one metric tensor more in the corresponding covariants. Thus, these relations can be used to recursively express all 4-point coefficients up to rank P (including the scalar integral) in terms of the coefficient $D_{0\dots 0}$ of rank $2P$ and 3-point integrals. For rank 4 or higher the coefficient $D_{0\dots 0}$ has the property that its Feynman-parameter integral involves a logarithmic integrand without any denominator and is, thus, well suited for numerical integration.

Figure 1 illustrates the reliability of the seminumerical method for an example where the Gram determinant $|Z|$ becomes small ($x \rightarrow 0$) and therefore the PV algorithm breaks down. Away from the tricky region, i.e. for increasing x (and thus $|Z|$), there is a transition region in which PV reduction and the seminumerical method agree within good accuracy until the latter becomes less precise than the PV method. This is due to the error of the numerically evaluated coefficient $D_{0\dots 0}$ which enters with a prefactor $|Z|$, and is thus suppressed for small $|Z|$.

The seminumerical method is limited to cases in which $|X|$ is not too small. If necessary, the case of both $|Z|$ and $|X|$ small, where PV reduction does not work either, can be cured with an expansion described below. For 3-point integrals, the soft or collinear singular cases have the property that $|X| = 0$ exactly, so that the seminumerical method is not applicable. However, these special cases are simple enough to be dealt with analytically as described in App. B of Ref. [2].

3.2. Expansion for small $|Z|$

The PV relations can be rewritten as

$$\tilde{X}_{0j} D_{i_1 \dots i_P} = 2 \sum_{r=1}^P \tilde{Z}_{ji_r} D_{00i_1 \dots \hat{i}_r \dots i_P}$$




Figure 1. A typical example for 4-point integrals with small $|Z|$ ($x \rightarrow 0$). The full diagram and the relevant subdiagram are given above; absolute predictions (in arbitrary units) for some tensor coefficients, relative deviations from PV reduction, and the kinematic specifications are shown below.




Figure 2. An example for 4-point integrals with both $|Z|$ and $|X|$ small ($x \rightarrow 0$). Details as in Fig. 1.

$$\begin{aligned}
& + |Z| D_{ji_1 \dots i_P} + 3\text{-point integrals}, \\
\tilde{Z}_{kl} D_{00i_1 \dots i_P} & \propto \sum_{n,m=1}^{N-1} \tilde{\tilde{Z}}_{(kn)(lm)} \left[f_n f_m D_{i_1 \dots i_P} \right. \\
& + 2 \sum_{r=1}^P (f_n \delta_{mi_r} + f_m \delta_{ni_r}) D_{00i_1 \dots \hat{i}_r \dots i_P} \\
& + 4 \sum_{\substack{r,s=1 \\ r \neq s}}^P \delta_{ni_r} \delta_{mi_s} D_{0000i_1 \dots \hat{i}_r \dots \hat{i}_s \dots i_P} \left. \right] \\
& + 2m_0^2 \tilde{Z}_{kl} D_{i_1 \dots i_P} - |Z| D_{kli_1 \dots i_P} \\
& + 3\text{-point integrals}, \tag{7}
\end{aligned}$$

which express the coefficients $D_{i_1 \dots i_P}$ and $D_{00i_1 \dots i_P}$ in terms of lower-rank 4- and 3-point coefficients and higher-rank 4-point coefficients dressed with a prefactor $|Z|$. For small $|Z|$, these relations can, thus, be used for an iterative determination of all 4-point coefficients from 3-point coefficients up to terms that are suppressed by some powers of $|Z|$. For a specific coefficient the iteration can be made arbitrarily precise upon including higher and higher ranks in the iteration.

Figure 1 demonstrates the reliability of the method for the same configuration as for the seminumerical method of the previous section. For increasing $|Z|$ (and x) the expansion becomes less and less precise, because the missing terms that are suppressed by some power of $|Z|$ grow. Again there is a region where all methods yield decent results.

The discussed expansion is limited to the case where $\tilde{X}_{0j}$ and $\tilde{Z}_{kl}$ are not too small for at least one set of indices j, k, l . If all $\tilde{X}_{0j}$ are small, then $|X|$ is small, too. This case is considered in the next subsection. The case in which all $\tilde{Z}_{kl}$ are small is further elaborated in Ref. [2].

3.3. Expansion for small $|Z|$ and $|X|$

The PV relations can again be rewritten as

$$\begin{aligned}
\sum_{r=1}^P \tilde{Z}_{ki_r} D_{00i_1 \dots \hat{i}_r \dots i_P} & \propto \tilde{X}_{k0} D_{i_1 \dots i_P} - |Z| D_{ki_1 \dots i_P} \\
& + 3\text{-point integrals}, \\
\tilde{X}_{ij} D_{i_1 \dots i_P} & = \text{const.} \times \tilde{Z}_{ij} D_{00i_1 \dots i_P} \\
& - 2 \sum_{r=1}^P \sum_{n=1}^{N-1} \tilde{\tilde{Z}}_{(in)(ji_r)} f_n D_{00i_1 \dots \hat{i}_r \dots i_P}
\end{aligned}$$

$$+ \tilde{X}_{0j} D_{ii_1 \dots i_P} + 3\text{-point integrals}, \tag{8}$$

which expresses the coefficients $D_{i_1 \dots i_P}$ and $D_{00i_1 \dots i_P}$ in terms of lower-rank 4- and 3-point coefficients and higher-rank 4-point coefficients dressed with a prefactor $|Z|$ or $\tilde{X}_{0j}$. If $|Z|$ and all $\tilde{X}_{0j}$ are small, these relations can be used to iteratively determine all 4-point from 3-point coefficients up to suppressed terms. Compared to the case of the previous subsection, where only $|Z|$ was considered small, the tensor rank grows faster with each iteration.

Figure 2 shows the reliability of the method for a specific example, revealing again the breakdown of PV reduction for sufficiently small x (and thus $|Z|$ and $|X|$), while the expansion becomes less precise for increasing x .

The expansion method fails if either all $\tilde{Z}_{ki_r}$ or all $\tilde{X}_{ij}$ are small. Possible treatments of these exceptional cases are also described in Ref. [2].

REFERENCES

1. G. Passarino and M. J. Veltman, Nucl. Phys. B **160** (1979) 151.
2. A. Denner and S. Dittmaier, Nucl. Phys. B **734** (2006) 62 [hep-ph/0509141].
3. Standard Model working group report of the workshop “Physics at TeV Colliders”, Les Houches, May 2005.
4. A. Denner, S. Dittmaier, M. Roth and L. H. Wieders, Phys. Lett. B **612** (2005) 223 [hep-ph/0502063] and Nucl. Phys. B **724** (2005) 247 [hep-ph/0505042].
5. A. Denner and S. Dittmaier, Nucl. Phys. B **658** (2003) 175 [hep-ph/0212259].
6. A. Ferroglia, M. Passera, G. Passarino and S. Uccirati, Nucl. Phys. B **650** (2003) 162 [hep-ph/0209219].
7. W. Giele, E. W. N. Glover and G. Zanderighi, Nucl. Phys. Proc. Suppl. **135** (2004) 275 [hep-ph/0407016]; R. K. Ellis, W. T. Giele and G. Zanderighi, hep-ph/0508308.
8. T. Binoth, J. P. Guillet, G. Heinrich, E. Pilon and C. Schubert, hep-ph/0504267.
9. S. Dittmaier, Nucl. Phys. B **675** (2003) 447 [hep-ph/0308246].
10. D. B. Melrose, *Nuovo Cimento* **XL A** (1965) 181.